Navigating Newborn Male Circumcision: Evidence, Practice, and Parental Counseling

Clinical Briefing and Role-Play Workshop



Facilitated by:
Ruth Mielke, PhD, CNM, FACNM, WHNP-BC

Gemini Notebook
Gemini Notebook

Learning Objectives


By the end of the presentation the learner will:

1. Describe the religious, cultural, and health reasons that parents choose newborn male circumcision
2. Describe the procedure of male newborn circumcision and the aftercare for circumcised newborns
3. Role play a conversation with parents who are considering newborn male circumcision

Gemini Notebook

What is male circumcision?


- Surgical removal of the foreskin (prepuce) covering the glans (head) of the penis.
- It exposes the glans and is typically performed during the newborn period, but can also be done later in life.



Gemini Notebook

The Epidemiological Landscape of Circumcision in the United States


National Incidence Trends



In-hospital newborn male circumcision incidence ranges between 55% and 59% (declining slightly from a peak of ~63% in 1999).

Note: These figures represent hospital discharges and may underestimate total rates by excluding out-of-hospital religious ceremonies.

Demographic and Regional Variance



- Geography: Incidence is highest in the Midwest (74%) and Northeast (67%), and lowest in the West (30%).
- Race/Ethnicity: Self-reported prevalence is highest among non-Hispanic White men (88%), followed by non-Hispanic Black men (73%), and Mexican-American men (42%).
- Socioeconomics: Circumcised newborns are disproportionately more likely to be privately insured than publicly insured, revealing a financial barrier to parental choice in states where Medicaid excludes the procedure.

Gemini Notebook

The Decision Triad: Why Parents Choose Circumcision


Medical & Health Drivers

Prevention of Urinary Tract Infections (UTIs), especially in the first year of life.

Reduced lifetime risk of acquiring HIV and other Sexually Transmitted Infections (STIs).

Prevention of penile cancer and phimosis (inability to retract foreskin).

Perception of easier genital hygiene.



Social & Cultural Drivers

Family tradition: The desire for the child to look like all the other men in the family (specifically the father).

Social conformity: The fear that the child will feel different from peers in locker rooms.

Clinical Context: Studies show up to 80% of parents make their decision before ever discussing it with a clinician, heavily relying on these social drivers.

Religious Drivers

Judaism (Brit Milah): An obligatory commandment performed on the eighth day of life by a mohel.

Islam (Tahara): Represents purification; can be performed in the neonatal period or as a rite of puberty

Gemini Notebook

The Health Impact Matrix: Weighing the Medical Evidence




Disease Category	Baseline Profile & Efficacy	Clinical Impact & Context
Urinary Tract Infections (UTIs)	**Baseline Risk (Uncircumcised):** ~1 in 100 chance in the first year of life. **Circumcised Risk:** ~1 in 1,000 chance.	A 10-fold reduction. Number Needed to Treat (NNT) is ~100 for normal infants, but drops to ~11 for infants with vesicoureteral reflux (VUR).
HIV Acquisition (Heterosexual)	40% to 60% risk reduction among heterosexual males in high-prevalence areas.	CDC models estimate a 15.7% lifetime risk reduction for all US males.
Other Sexually Transmitted Infections	**HPV:** 30% to 40% reduction in risk. **HSV-2:** 28% to 34% reduction in risk.	HPV reduction protects against genital warts and reduces transmission of high-risk cervical cancer strains to female partners. **Syphilis** demonstrates a protective trend.
Penile Cancer	Uncircumcised men face an Odds Ratio (OR) of 2.3 for invasive squamous cell carcinoma.	Absolute incidence remains exceedingly rare in developed nations.

Gemini Notebook

Evaluating Procedural Complications and Surgical Risks

Acute Complications (Newborn Period)

- Incidence: Significant acute complications are rare, occurring in approximately 0.19% to 0.22% (roughly 1 in 500) of newborn procedures in US hospitals.
- Primary Risks: Minor bleeding (most common, easily controlled with pressure or silver nitrate), local infection, and imperfect tissue removal.



The Risk Multipliers

- Timing: Complication rates rise exponentially if the procedure is deferred beyond the newborn period, eventually requiring general anesthesia.
- Provider: Untrained providers yield vastly higher complication rates than medical professionals.
- Infection Warning: The ultra-Orthodox practice of metzitzah b'peh (direct oral suction) poses a severe risk of neonatal Herpes Simplex Virus (HSV) transmission and should be strictly counseled against.

Late Complications

- Manifestations: Penile adhesions, vascularized skin bridges, redundant prepuce, and meatal stenosis.
- Prematurity Risk: Circumcising premature infants or those with prominent suprapubic fat pads carries a higher risk of poor cosmesis or trapped penis.

Gemini Notebook


Procedural Modalities and Clinical Contraindications

Core Surgical Principles


1. Estimation of external skin removal.
2. Dilation of the preputial orifice and visual inspection of the glans.
3. Blunt lysis of physiologic adhesions.
4. Device placement for hemostasis, followed by surgical excision.



Gomco Clamp: Utilizes a bell and base mechanism; left in place for 3-5 minutes for hemostasis before scalpel excision.



Plastibell Device: A plastic ring inserted under the foreskin with a suture tie; the necrotic tissue and ring fall off spontaneously in 5-8 days.



Mogen Clamp: Two flat blades with a 3mm slit that crush the skin to create hemostasis prior to excision.




Absolute Contraindications

Instability: Prematurity or active illness.
Anatomy: Congenital anomalies (hypospadias, congenital chordee, penoscrotal webbing, buried penis).
Hematology: Family history of bleeding disorders or lack of routine Vitamin K administration.

Gemini Notebook

The Analgesia Pyramid: Mandating Effective Pain Control

Newborn circumcision causes significant pain, evidenced by altered heart rate, oxygen saturation, and crying. The AAP mandates adequate analgesia.



Apex: **Regional Nerve Blocks** (Most Effective)

- **Dorsal Penile Nerve Block (DPNB):** Bilateral injections of 1% lidocaine (without epinephrine) at the base of the penis. Highly effective, superior to topicals.
- **Subcutaneous Ring Block:** Circumferential injection at the base/midshaft. Highly effective but has a ~5-8% failure rate.

Middle: **Topical Anesthetics (Pharmacologic)**

- 4% Lidocaine (LMX4): Faster onset (30 mins); preferred over EMLA as it carries no risk of methemoglobinemia.
- EMLA Cream: Requires 60-90 minutes of application under a plastic wrap occlusion.
- **Warning:** Topicals may cause blistering in low-birth-weight infants; nerve blocks are preferred for this population.

Base: **Non-Pharmacologic Adjuncts** (Insufficient Alone)

- **Methods:** Sucrose pacifiers, swaddling, physiologic positioning in a padded environment.
- **Rule:** These modify the environment and soothe the infant but do not block surgical pain pathways.

Gemini Notebook

The Aftercare Dichotomy: Parental Discharge Instructions



Care of the Circumcised Penis (Acute Wound Healing)

- **Immediate Care**: Apply petroleum jelly (Vaseline) to the bandage with every diaper change to prevent tissue from sticking. (Note: If Plastibell is used, do not pull the ring; it falls off in 5-8 days).
- **Hygiene**: Sponge baths only until fully healed (7-10 days). Wash off stool gently; do not wipe the glans.
- **Normal Progression**: The glans will appear red, then form a yellowish film (granulation tissue). This is normal healing, not pus.
- **Red Flags (Call Provider)**: Active bleeding (beyond minor spotting), no urine output within 12 hours post-procedure, or persistent swelling/fever.



Care of the Uncircumcised Penis (Physiologic Normalization)

- **Immediate Care**: Wash the outside of the penis gently with mild soap and water, just like the rest of the body.
- **The Golden Rule**: DO NOT FORCIBLY RETRACT THE FORESKIN.
- **Physiological Progression**: In 96% of newborns, the foreskin is naturally fused to the glans. It will separate on its own timeline over the **first 3 to 10 years of life** as smegma accumulates and facilitates separation.
- **Hygiene**: Once natural retraction occurs years later, the child can be taught to retract, clean, and replace the foreskin.

Gemini Notebook

The Ethical Framework of Informed Consent

Surrogate Decision-Making

Because infants cannot provide consent, parents act as surrogate decision-makers. The AAP dictates that parents are granted wide latitude to make decisions based on their child's best interest, which legally and ethically includes the family's cultural, religious, and ethnic traditions.



The Clinician's Mandate

Objective and Nonbiased: The clinician's duty is to explain potential benefits and risks objectively, ensuring parents understand circumcision is an elective procedure, not a medical mandate.

Timing: Counseling should ideally occur before conception or early in pregnancy.

The Compass, Not the Steering Wheel

Health Literacy: The informed consent process historically falls below the standard of care. Avoid medical jargon. Your role is not to dictate the path. It is to provide a reliable clinical compass—orienting the parents to the medical realities so they can navigate the decision according to their own family values.

Gemini Notebook

The Counseling Flowchart: Mapping the Conversation

Step 1: Assess Values & Context

Action: Ask open-ended questions to discover the parents' baseline knowledge and motivations.

Phrasing: Have you and your partner discussed your thoughts on circumcision yet? What factors are most important to you?

Step 2: State Objective Facts (Risks & Benefits)

Action: Distill the Evidence Matrix and Complication Scale without leading the parent.

Phrasing: Medically, the procedure lowers the risk of UTIs in the first year and offers protection against certain STIs later in life. However, it is surgery, and carries a small risk of bleeding or infection.

Step 3: Address Pain Control & Procedure

Action: Reassure parents regarding the AAP's mandate for analgesia.

Phrasing: If you choose this route, we use highly effective local nerve blocks and comforting techniques so your baby is as comfortable as possible.

Step 4: Teach-Back & Support Decision

Action: Confirm understanding of aftercare (for whichever path they choose) and validate their autonomy.

Phrasing: To make sure I explained the aftercare clearly, can you tell me how you'll handle his diaper changes next week? Whatever you decide, we fully support your choice.

Gemini Notebook

Clinical Workshop: Role-Play Scenario

The Setup

You are conducting a 34-week prenatal visit. The parents, Ana and Jamal, bring up the topic of circumcision for their expected son.

Jamal (Leaning Yes): All the men in my family had it done. I just don't want him to look different or get teased in the locker room. Plus, isn't it cleaner?

Ana (Leaning No): I've been reading online that it's a traumatic mutilation and the baby feels everything. Why would we put him through surgery if it isn't medically broken?



The Prompt:

They look at you and ask, "You're the expert. What do you think we should do?"

Your Objective:

Break into groups of three (Home Visitor, Ana, Jamal). Utilize the Counseling Flowchart to:

1. Validate both of their concerns without taking a side.
2. Correct the misinformation regarding pain control and hygiene.
3. Guide them to consider their own values (religious, social) as they consider medical data.
4. Guide them to evidence based resources that will address questions that you cannot answer.

Gemini Notebook

Insurance Coverage of Newborn Male Circumcision

- **Insurance coverage is not uniform across all insurance carriers.**
- **Some private carriers cover it but 17 states (California included) do not cover routine NMC under Medicaid.**
- **Medi-Cal Fee-for-Service** (FFS) system covers about 13.3% of Medi-Cal beneficiaries. Under the FFS delivery plan a circumcision is covered only if the provider deems it medically necessary. CPT codes 54150 and 54160 are non-benefits, but may be available for medical necessity with prior authorization. **Medi-Cal Managed care** is the vast majority of Medi-Cal beneficiaries - 86.7%. *A managed care plan has the option to cover other services in addition to the required FFS benefits. In other words, CHS administrators have chosen to cover neonatal circumcisions based on American Academy of Pediatrics recommendations.* Medi-Cal recipients covered by a managed care plan should check with their provider to ascertain the circumstances under which a circumcision procedure would be covered.

Medically necessary conditions may be: Phimosis (foreskin too tight and cannot be retracted), balanitis (inflammation of the glans) or chronic UTIs

https://biologyinsights.com/is-circumcision-covered-by-insurance-in-california/

Cost of Newborn Male Circumcision

- Total cost without insurance typically falls between $250 and $400 for the procedure itself, plus an additional $235 to $375 in hospital facility charges if it’s performed during the hospital stay.
  - That puts the full cost somewhere in the range of $485 to $775.
- Many hospitals offer self-pay discounts or payment plans for uninsured families.
- If your client is interested or planning NMC, call managed care plan/insurance before delivery and ask if procedure is included in newborn care.



REFERENCES/RESOURCES

1. Mielke, R. T. (2013). Counseling parents who are considering newborn male circumcision. Journal of Midwifery & Women's Health, 58(6), 671-682. https://doi.org/10.1111/jmwh.12130
2. O'Connor, S. D. (2023). Making Choices—Newborn Male Circumcision. Narrative inquiry in bioethics, 13(2), 107-111. https://doi.org/10.1353/nib.2023.a909671
3. American Academy of Pediatrics (10/10/2025) Healthychildren.org. Circumcision: A Parent's Choice.
4. Google Gemini Notebook. (October, 2026) Infographic: Newborn Male Circumcision: Weighing Benefits, Risks, and Health Outcomes.
5. PedsDocTalk(2026, November 2). Circumcision: The Medical Facts Without the Pressure [Video]. YouTube.https://youtu.be/v27oN7dAVP8.